

	Department Name Address	ALS	Paramedic
		Revision #	
		Implementation Date	
Protocol	2.2.2 Airway Management- Pediatric	Last Reviewed/Update Date	
Author / Owner		Medical Director	

Treatment goals in airway management of a patient in the pre-hospital environment include ensuring adequate oxygenation and safe, timely transport to the appropriate care facility.

1. Baseline care standards.
2. Place patient in position of comfort and reassure.
3. Maintain cervical spine control on patients with suspected trauma.
4. Auscultate with stethoscope at least four different areas of the chest and document.
5. If spontaneous breathing is present without compromise:
 - Monitor breathing during transport.
 - Administer oxygen PRN:
 - Infants via infant mask @ 2 – 4 L per minute
 - Small child (1 – 8 years) via pediatric mask @ 6 – 8 L per minute.
 - Older child (9 – 15 years) via non-rebreather @ 15 L per minute.
 - If mask is not tolerated, administer via blow-by method.
6. If spontaneous breathing is present with compromise:
 - Manually open airway.
 - Suction PRN.
 - Administer oxygen PRN:
 - Infants via infant mask @ 2 – 4 L per minute
 - Small child (1 – 8 years) via pediatric mask @ 6 – 8 L per minute.
 - Older child (9 – 15 years) via non-rebreather @ 15 L per minute.
 - If mask is not tolerated, administer via blow-by method.
 - If unable to maintain airway, insert oropharyngeal or nasopharyngeal airway PRN.
 - Assist ventilations with BVM.
 - Monitor O₂ saturations with pulse oximeter.
7. If spontaneous breathing is absent or markedly compromised:

- Manually open airway.
 - Suction PRN.
 - If unable to maintain airway, insert oropharyngeal or nasopharyngeal airway.
 - Ventilate with BVM @ 20 per minute for a child and 30 per minute for an infant.
 - Monitor O₂ saturations with pulse oximeter.
8. If patient is unresponsive and unable to maintain an adequate airway or without a pulse, intubate using direct laryngoscopy and appropriate size endotracheal tube. If cervical spine injury is suspected, have second person maintain c-spine control during intubation.
- Verify tube placement by auscultation of breath sounds, esophageal detector device, and capnometry.
 - Secure ETT with commercial device.
 - Monitor O₂ saturations with pulse oximeter.
9. Insert NG tube to decompress stomach PRN.
10. Consider needle cricothyrotomy if indicated.

Medical Director's Signature

Date

Disclaimer:

The protocols have been developed by the North Dakota Department of Health are meant to be used as general guidance for developing protocols for individual emergency medical services agencies. These sample protocols are not meant to be medical or legal advice; nor do they establish standards of care. Each emergency medical services agency must tailor protocols based on their specific needs or capabilities. Local medical directors must be consulted with and approve any protocol(s) prior to becoming operational in an emergency medical services agency. The North Dakota Department of Health make no representation on the accuracy of information contained herein and accepts no liability for any loss or damage arising from any content error or omission.